



Breaking the 1 m s^{-1} Barrier

Data Collection & Data Analysis

Arpita Roy



Future of the Field

Heralding **Extreme Precision** Spectroscopy

Instruments attempting to achieve $\sim 10\text{cm/s}$ precision for the detection of Earths




Basic Philosophy

Stabilize everything you can in hardware

Build in diagnostics for other variables so you can correct in software


Instrument Design

Maximize Stability Optically + Thermo-mechanically



Want instrument to deliver high-fidelity spectra so that you can

- Trust your line profiles
- Interpret variations as astrophysical phenomenon rather than instrument + Earth systematics
- Not directly applicable to iodine instruments [Need extra processing/forward modeling]




Data Pipeline

Need **extreme**
precision analysis to
produce extreme
precision RVs!


Radial Velocity Measurement Technique

Cross Correlation with Mask or Template

True bulk motion causes all stellar lines to move in an identical way [without changing shape]



Measuring Radial Velocity Via Gaussian Fitting



Pick best areas, rich in RV information and clean from tellurics

Works well for FGK stars with well defined lines and continuum

Mask needs to evolve over time as the data stream lengthens

Very hard for M dwarfs: lack of clean lines + poor synthetic models

**Let's Get To Know +
[♥] Our Data**


At the Start of Analysis

Beautiful (albeit complex) CCD images

All your careful instrument design effort is already evident in these images



- Well focused spots
- Minimal scattered light
- Low aberrations
- No bright ghosts
- Minimal slit tilt
- Many more...




Habitable Zone Planet Finder: Raw frame of Barnard's Star with laser frequency comb calibration

At the Start of Analysis

Beautiful (albeit complex) CCD images

All your careful instrument design effort is already evident in these images – **OR NOT!**




SOPHIE early data

Basic++

Image Processing


Introduction to RV Data Analysis

Image Processing




Things to be careful about:

- Bad pixels (change over time)
- Flat edges
- Pixel-to-pixel variation



Raw Image



Bias



Flat



Introduction to RV Data Analysis

Image Processing

Aside: Detectors are not perfect!

Example A: Pixel Size Inhomogeneity

Adds noise to the spectrum and causes errors in wavelength calibration



Correction: Flux effect should be removed with flat fielding

Wavelength error might be possible to correct if every sub-pixel shape mapped out in laboratory


Introduction to RV Data Analysis

Image Processing

Aside: Detectors are not perfect!

Example B: Charge Transfer Inefficiency

Changes continuum level and skews line shape in direction of readout



Correction: Measure CTI for your CCD, consider correcting in pipeline
Difficult because flux dependent - need library of flats at different SNR


Introduction to RV Data Analysis

Image Processing

Aside: Detectors are not perfect!

Example C: Fringing

Makes it difficult to use redder wavelengths on detector – important for M dwarfs!




Correction: Contemporaneous flats not practical
Need to solve this in hardware with optimized CCD coatings

Basic++

Data Reduction


Introduction to RV Data Analysis

Spectral Extraction



Things to be careful about:


- Curved order rectification
- Slit tilt
- Cross-talk between orders
- Scattered light/ghosts
- Stitching boundaries



Order Trace





Remove cosmic rays and bleeding from contiguous orders



Optimally Extract Spectra

Introduction to RV Data Analysis

Wavelength Calibration




Things to be careful about:

- Emission lamps have differential drift between species, aging effects, contaminants
- Fabry-Perot etalons must be evaluated for drift (including chromatic drift)
- Laser frequency combs are so nice and narrow that they can cause sampling problems

Instrument-Based Corrections

Introduction to RV Data Analysis

Flux Calibration




Things to be careful about:

- Overall continuum fluctuates a lot based on clouds/seeing/guiding
 - Scrambling cannot restore light outside fiber
 - Light loss is chromatic (based on level of atmospheric dispersion correction)

Introduction to RV Data Analysis

Flux Calibration




Things to be careful about:

- Blaze curve within an order varies depending on echelle illumination (variable with guiding, seeing etc.)
 - Reweights lines and can skew line shapes
 - Want to remove variability but not inherent photon noise information

Introduction to RV Data Analysis

Instrument Drift Correction

Fiber-fiber drift is our main diagnostic of instrument stability




Introduction to RV Data Analysis

Instrument Drift Correction

Keep an eye on absolute fiber drift for systematic changes


Especially daily, seasonal, or yearly cycles



Earth-Based Corrections

Introduction to RV Data Analysis

Barycentric Correction



Need to calculate barycentric motion at flux-weighted mid-point of observation

[Cannot assume exposure was uniformly illuminated – variables include guiding, seeing, airmass, clouds etc.]

Need chromatic barycentric correction


[Different color photons arrive at different rates]

Introduction to RV Data Analysis

Sky Correction

a. Telluric Emission

Sky emission lines contaminating the activity-sensitive sodium D absorption lines



Solution: Subtract sky fiber spectrum (might need to be scaled for slightly different fiber)



Seems to work well with even simple subtraction

Introduction to RV Data Analysis

Sky Correction

b. Telluric Absorption


Left untreated, telluric lines will have very large impacts on RV precision.



Introduction to RV Data Analysis

Sky Correction

b. Telluric Absorption



Micro-telluric lines (depth of <1%) are particularly insidious because they are everywhere in the spectrum and cannot be effectively excluded from the CCF mask

Cunha et al. (2014) showed they can impact RVs at the 20 cm/s level or greater if not carefully treated.


Solution: Forward modeling based on sky fiber + measured telluric spectrum * at telescope site * over different seasons

Introduction to RV Data Analysis

Sky Correction

c. Scattered Sunlight Contamination

In general, spectral contamination is a concern
Temporal variability is usually present due to barycentric + stellar motion



Possible Culprits Include:

Other stars, both bound and background

Faint close-by objects - level of contamination is seeing + orientation dependent

Moonlight

Zodiacal Light





Unexpected sources must be diagnosed directly from the data [eg. Wright, Roy et al. 2013]

Introduction to RV Data Analysis

Sky Correction

c. Scattered Sunlight Contamination



Sunlight is particularly insidious when observing Sun-like stars since the contaminating spectrum is a good match to the mask lines

Solution: Sky fiber is essential

- Correct directly from sky fiber in CCF space, or
- Model sky spectrum and correct in spectral space
- Can also replace sky fiber with broadband proxy for faint sky


Star-Based Corrections

Stellar Activity

How Do We Deal With It?

Capture as many activity indicators in bandpass as possible



This also maximizes Doppler information content




Stellar Activity

How Do We Deal With It?

KPF Green Channel



KPF Red Channel




+ Separate Ca H&K spectrophotometer

Stellar Activity

How Do We Deal With It?

Measure Known Activity-Sensitive Lines



Start with Canonical Activity Indicators

Eg. Ca H&K from Isaacson & Fischer 2010


Mainly interested in relative variability but good to have handle on absolute activity levels as well

Stellar Activity

How Do We Deal With It?

Measure Known Activity-Sensitive Lines

Check for correlations with RV



Important Lessons

Stellar activity not coherent over long timescales

Need tight cadence over a period of time to see this


Stellar Activity

How Do We Deal With It?

Scrutinize Line Profiles: Bisectors & FWHM

If the instrument profile is stable then any line variations can be ascribed an astrophysical origin

Bisectors have long been used to distinguish bulk motion of the star from spectral line asymmetries due to activity or contamination




- True Doppler motion should not affect line shapes
- Single lines have low SNR in high resolution spectroscopy
- Use the cross correlation function as an "aggregate line"


Stellar Activity

How Do We Deal With It?

Scrutinize Line Profiles: Bisectors & FWHM



Measuring Radial Velocity Via Gaussian Fitting of Cross Correlation Function (CCF)





Bisector variation is caused by both stellar activity and spectral contamination

Need to be very careful which lines to "average"
Need to be very careful what wavelength range to use
[otherwise dilute chromatic effect]

Stellar Activity

How Do We Deal With It?

Scrutinize Line Profiles: Bisectors & FWHM



Example: Marvells-1b

Suspicious planet signal

Investigation revealed extremely variable bisectors



[Took a while to see this because of lack of stabilized instruments]

$$\text{Bisector Inverse Slope [BIS]} = v_{top} - v_{bottom}$$

Stellar Activity

How Do We Deal With It?

Scrutinize Line Profiles: Bisectors & FWHM



Example: *Marvels-1 b*

Clear correlation between RV and both BIS and FWHM
Modeling showed it was a rare face-on binary masquerading as a double planet system

Unfortunately stellar activity signal not so coherent or clean

Observing Strategy


Cadence + Total Time

Necessary in Addition to Instrument Precision

To End On a Positive Note

Looking good to $\sim 30\text{cm/s}$, with path to 10cm/s

Pipelines perform well on quiet standard stars [with best instruments so far]



New State of the Art Being Established Right Now*:
In the optical, ESPRESSO getting to $\sim 30 \text{ cm/s}$ over few days
In the NIR, HPF at $\sim 1.3 \text{ m/s}$ over 2 weeks

* on sky during commissioning